



County of Erie

CHRIS COLLINS
COUNTY EXECUTIVE

DEPARTMENT OF HEALTH

ANTHONY J. BILLITTIER IV, M.D., FACEP
COMMISSIONER OF HEALTH

HEALTH UPDATE #248 ANNUAL LYME DISEASE AND LEPTOSPIROSIS REPORT

JULY 9, 2009

Please distribute to Emergency Departments, Infection Control Departments, Infectious Disease Departments, Pediatrics, Director of Nursing, Medical Director, Laboratory Director, all patient care areas, Veterinarians, Animal Control Officers, and Wildlife Rehabilitation Specialists.

Attached are two reports based on the 2008 Lyme Disease and Leptospirosis Veterinarian Survey conducted by the Vector Control Program of the Erie County Department of Health (ECDOH). The Lyme disease report also includes human statistics from the Epidemiology Program of the ECDOH as well as Tick identification data from both the New York State Department of Health (NYSDOH) and the Erie County Vector Lab. Each of these reports has epidemiologic and zoonotic importance.

Please direct any questions regarding these reports to John Eiss or Peter Tripi at the ECDOH, Vector Control Program at 716-961-7524.

Attachments:

Lyme 2005 – 2008

Leptospirosis 2004 - 2008

Health Category Definitions:

Health Alert FLASH: conveys the highest level of importance due to a large-scale, catastrophic public health emergency; warrants immediate action or attention

Health Alert Priority: conveys the highest level of importance; warrants immediate action or attention to a health problem or situation

Health Advisory: provides important information for a specific incident or situation; may not require immediate action

Health Update: provides updated information regarding an incident or situation; no immediate action necessary



**Erie County Health Department
Vector Surveillance & Control Field Laboratory
Buffalo, New York**



2005 - 2008 Lyme Disease Survey

6/ 2009

Evidence suggests Lyme disease can be acquired by humans and dogs locally in Erie County.

Lyme disease is the most prevalent arthropod-borne disease in New York State, and has been endemic in the lower Hudson Valley and Long Island since at least the 1980s. Evidence suggests that Lyme disease is expanding into the Western Region of the State, including Erie County.

Number of Human Lyme Disease Cases Confirmed by the NYSDOH in Erie County							
2008	2007	2006	2005	2004	2003	2002	2001
7*	7*	4	11	1	1	1	4



***Note: Prior to 2007, travel history outside Erie County was not routinely obtained. In two of the 2007 and in three of the 2008 cases there was no significant travel history outside Erie County reported.**

The Erie County Health Department Vector Field Laboratory has provided a regional tick identification service since 2001. The ability to identify ticks and determine the degree of engorgement in a timely manner can be a valuable tool in early diagnosis and treatment of Lyme disease.

Ticks Identified in Erie County*			
Year	Total Ticks Identified	<i>Ixodes scapularis</i>	% <i>Ixodes scapularis</i>
2008	265	92	34.7%
2007	184	62	33.7%
2006	164	32	19.5%
2005	76	21	27.6%
2004	49	8	16.3%
2003	84	33	39.3%
2002	103	16	15.5%
2001	49	9	18.4%
Total	974	273	28.0%

*Note: Ticks were identified by the Erie County Vector Field Laboratory, NYSDOH Western Regional Entomologist and the NYSDOH Hudson Valley Community College.





With respect to Lyme disease, dogs may be considered sentinel animals because they are at greater risk of tick infestation than humans. They are compliant and easily sampled, and have a pronounced antibody response to the spirochete infectious agent. Moreover, since dogs frequently develop asymptomatic disease that can lead to lameness, their owners are often motivated to have their animals tested.

Erie County Canine Lyme Disease Survey Results				
Year	Vets Surveyed	Vets That Test for Lyme	Canines Positive^o	No Travel History
2008	33	25	63*	11
2007	47	34	74	19
2006	54	35	83	23
2005	66	47	54	22
Total	200	141	274	75

^oNote: Laboratory confirmation of Lyme disease relies on indirect methods such as antibody detection. Scientific studies have demonstrated variable levels of sensitivity (30-80%) and specificity (80-90%) associated with the performance of these assays and are dependant upon stage of the infection, presence of cross-reacting antibodies, etc. Therefore, it must be assumed that a laboratory result for the diagnosis is not absolute and performance characteristics of the assay must be considered. Additionally, the diagnosis of the disease cannot rely on the laboratory result alone, but must incorporate clinical recognition, history, and other pertinent information.

*One of the 63 positive cases in 2008 was a cat.

Discussion: In 2008 and 2007, five confirmed human cases of Lyme disease reported no travel history outside Erie County. A large percentage of the ticks identified in Erie County are *Ixodes scapularis* (Blacklegged deer tick), the vector of Lyme disease. Substantial numbers of confirmed canine Lyme disease cases with no travel history have occurred over the last four years.

Conclusion: Evidence suggests Lyme disease can be acquired by humans and dogs in Erie County. Lyme disease should be considered a Public Health threat in Erie County.



For information contact:

John P. Eiss john.eiss@erie.gov
Public Health Consultant

or

Peter Tripi peter.tripi@erie.gov
Senior Public Health Sanitarian

or

Glenn Robert robertg@erie.gov
Senior Investigating Sanitarian

Vector Surveillance and Control Program



Tick Identification Service:
Erie County Health Department
Vector Surveillance & Control Program
462 Grider Street Room BB-122
Buffalo, NY 14215
716-961-7524



**Erie County Health Department
Vector Surveillance & Control Field Laboratory
Buffalo, New York**

**Leptospirosis
2004-2008 Survey and Report**

06/2009

The use of covered garbage totes may have contributed to the drop in canine leptospirosis cases in Erie County over the past five years.

Overview:

Leptospirosis is a bacterial disease associated with wild and domestic animals. Many different kinds of animals carry the bacterium; while some may become sick many will have no symptoms. Leptospira organisms have been found in cattle, pigs, horses, dogs, rodents, and wild animals. The disease has a seasonal incidence during the late spring, summer and fall, when the soil is moist and alkaline. Rainfall and higher temps seem to increase the number of cases in dogs.

Animals and humans become infected through contact with water, food, or soil contaminated by urine from these infected animals. This may happen by swallowing contaminated food or water or through skin contact, especially with mucosal surfaces, such as the eyes or nose, or with broken skin. The disease is not known to be spread from person to person.


In humans, leptospirosis can cause a wide range of symptoms and some infected persons may have no symptoms at all. Symptoms include high fever, severe headache, chills, muscle aches, and vomiting, and may include jaundice (yellow skin and eyes), red eyes, abdominal pain, diarrhea, or a rash. Many of these symptoms can be mistaken for other diseases. If the disease is not treated, the patient could develop kidney damage, meningitis (inflammation of the membrane around the brain and spinal cord), liver failure, and respiratory distress. In rare cases death occurs. The incubation period is usually 10 days with a range of 4 to 19 days. The disease is diagnosed using specific blood tests available through public health laboratories. The antibiotics of choice are penicillin, streptomycin, tetracycline and erythromycin. Kidney dialysis may be necessary in some cases.

Human:

“Epidemic leptospirosis most commonly occurs after flooding in densely populated centers in developing countries. Also, experience [has shown] there is a high rate of misdiagnosis of leptospirosis” (Spichler et. al.). According to the CDC the reported incidence of leptospirosis are 100–200 cases per year in the United States with most (50–100 cases) occurring outside the continental United States in Hawaii. Leptospirosis is likely under-diagnosed in the United States, with reported incidence depending largely upon clinical index of suspicion. New York State DOH reports 1 human leptospirosis case in 1994, 1996 and 1997 with 3 cases reported in 2000. More recent information is not available. None of the cases were in Erie County.

Animal:

Many articles and studies have been published in recent years indicating leptospirosis is a reemerging disease among dogs in the US and Canada. Cats are not known to be affected. In a paper that was published in Ontario, Canada, documenting a large increase of canine leptospirosis in 2000, the following was stated: “The reasons may be the increased and endemic infection of urban wildlife (notably raccoons, skunks) with leptospirosis, combined with increased numbers of urban wildlife and an increasing index of suspicion by veterinarians, thus promoting serological testing. Although canine leptospirosis is recognized to have been increasing in Ontario in the last few years, the fall of 2000 saw a marked rise in the number of cases. A major factor was probably the wet and exceptionally warm late summer and fall, which provided conditions that were ideal for the transmission of *Leptospira* from wildlife.” (Prescott et. al.)



leptospire organism

Results of the survey of small animal veterinary hospitals conducted by the Erie County Health Department in 2004-2008:

(The survey was conducted by phone and email.)

Leptospirosis Survey: Case Results					
	2004	2005	2006	2007	2008
Small Animal Vet Hospitals Participating	66	66	51	47	33
Canine Blood Samples Sent for Leptospirosis Testing	181	186	143	99	52
Cases Confirmed as Leptospirosis <small>(note 1 & 2)</small>	60	47	39	15	7
Suspected Leptospirosis Cases Unconfirmed <small>(note 3)</small>	14	20	31	19	16
Total Confirmed and Suspected Leptospirosis Cases	74	67	70	34	23

Note 1: The testing for leptospirosis was performed by six laboratories: Cornell, Antech, Idexx, Priority, MSU, Vitatech.


Note 2: Laboratory confirmation of leptospirosis relies on indirect methods such as antibody detection. Scientific studies have demonstrated variable levels of sensitivity (30-80%) and specificity (80-90%) associated with the performance of these assays and are dependant upon stage of the infection, presence of cross-reacting antibodies, etc. Therefore, it must be assumed that a laboratory result for the diagnosis is not absolute and performance characteristics of the assay must be considered. Additionally, the diagnosis of the disease cannot rely on the laboratory result alone, but must incorporate clinical recognition, history, and other pertinent information.

Note 3: In most cases if a dog was not tested it was because of a cost issue with the owner.

Leptospirosis Survey: Serovar Results (note 4)					
	2004	2005	2006	2007	2008
Grippe Serovar	10	8	11	4	2
Pomona Serovar	10	7	8	2	1
Ictero Serovar (associated with rats)	20	27	34	9	5
Canicola Serovar	3	5	8	2	1
Bratislava Serovar (associated with rats)	18	14	17	6	1
Autumnalis Serovar	18	12	11	8	1
Other Serovars Found	2	2	2	1	1

Note 4: Multiple serovars can be detected in a single positive leptospirosis serum sample

Number of Laboratory Confirmed Canine Leptospirosis Cases From 2004 to 2008 by Township



Note: Numbers based on locations of small animal veterinarian hospitals surveyed

Leptospirosis Survey: Veterinary Hospital Vaccination Results (note 5)

	2004	2005	2006	2007	2008
Hospitals That Vaccinate All Canines For Leptospirosis	33	19	20	17	9
Hospitals That Vaccinate On a Case By Case Basis	23	37	22	28	20
Hospitals That Do Not Vaccinate For Leptospirosis	10	10	9	2	4
Hospitals Participating in The Survey	66	66	51	47	33

Note 5: The following leptospirosis vaccines are used by Vets in Erie County: Duramune (Fort Dodge) and Vanguard L4 (Pfizer – Schering Plough).

Discussion:

National and local leptospirosis incidence data is not readily available due to a lack of reporting requirements. As a human illness, leptospirosis is probably under-diagnosed.


Confirmed cases of canine leptospirosis in Erie County have dropped every year of the survey. In 2004 there were 60 confirmed cases and in 2008 there were 7 confirmed cases. Increased vaccination of canines, better vaccines, changes in annual weather conditions and the number of veterinarians responding to the survey may account for some of the annual drop.

Another factor may be the large drop in the rat population in certain municipalities of Erie County as they employ covered garbage totes along with an aggressive rodent control program. During the years of the survey the municipalities with the highest number of rodent complaints has corresponded to the townships with highest number of canine leptospirosis cases. Evidence now indicates the drop in canine leptospirosis cases corresponds to the drop in rat populations in these same townships.

During the course of the survey there were 168 confirmed cases of leptospirosis with a total of 291 leptospirosis serovars detected. 57.5% of those serovars are associated with rats.

177 out of 279 leptospirosis serovars identified in the Erie County survey are included in currently available six month leptospirosis vaccines. Use of a canine leptospirosis vaccine may be indicated at least on a case-by-case basis. As canine leptospirosis appears to be more prevalent during the summer and fall in Erie County, it may be more important to protect dogs during this period.

Leptospirosis is infectious to humans; it is important for veterinary hospital personnel, Animal Control Officers, the SPCA wildlife rehabilitators and other wild animal handlers to take precautions to avoid possible infection. Wearing gloves and avoiding contact with animal urine are advisable.



For information or comment contact:
 Vector Surveillance & Control Program
 Erie County Health Department

John P. Eiss john.eiss@erie.gov
 Associate Public Health Sanitarian

Peter Tripi peter.tripi@erie.gov
 Senior Public Health Sanitarian

Glenn Robert robertg@erie.gov
 Senior Investigating Sanitarian