



COUNTY OF ERIE

MARK POLONCARZ

COUNTY EXECUTIVE

GALE R. BURSTEIN, MD, MPH, FAAP
COMMISSIONER OF HEALTH

DEPARTMENT OF HEALTH

HEALTH ADVISORY #349

December 13, 2018

MEASLES EXPOSURE IN ERIE COUNTY

Please distribute to the Chief Medical Officer, Infection Control Department, Infectious Disease Department, Pediatric Department, Director of Nursing, Emergency Department, Primary Care Clinics, and all patient care areas.

SUMMARY

- The Erie County Department of Health (ECDOH) has confirmed that a recent emigre treated at Millard Fillmore Suburban Hospital has been diagnosed with measles. This individual has visited multiple venues in Erie County potentially exposing others to measles between December 4, 2018 and December 11, 2018.
- Anyone who visited the following locations in Erie County may have been exposed to measles:
 - Catholic Charities, 20 Herkimer Street, Buffalo on 12/4/2018 between 11:00am and 5:00pm
 - Erie County Department of Health Clinic, 608 William Street, Buffalo on 12/4/2018 between 11:30am and 4:00pm
 - Sweet Home High School, 1909 Sweet Home Road, Amherst on 12/5/2018 between 5:00pm and 10:30pm
 - Erie County Department of Social Services, 158 Pearl Street, Buffalo on 12/6/2018 between 8:00am and 1:00pm
 - Erie County Department of Health Clinic, 608 William Street, Buffalo on 12/6/2018 between 10:30am and 2:00pm
 - Catholic Charities, 20 Herkimer Street, Buffalo on 12/6/2018 between 11:30am and 3:00pm
 - Erie County Department of Social Services, 158 Pearl Street, Buffalo on 12/6/2018 between 12:30pm and 4:30pm
 - ALDI, 4259 Transit Road, Buffalo on 12/6/2018 between 5:00pm and 9:00pm
 - Best Buy, 4401 Transit Road, Williamsville on 12/6/2018 between 6:00pm and 9:15pm

- Marshalls, 2383 Maple Road, Williamsville on 12/6/2018 between 6:00pm and 10:00pm
 - Millard Fillmore Suburban Hospital Emergency Department, 1540 Maple Road, Williamsville on 12/10/2018 between 9:00am and 5:30pm
- Individuals who were exposed and not immune to measles, as defined below, could develop signs and symptoms of measles 7-21 days after exposure.
 - Healthcare providers should have a high index of suspicion in patients who present with symptoms consistent with measles and have visited the locations listed. To expedite public health containment strategies, providers should implement appropriate infection control measures when measles is suspected and report immediately to the local health department (LHD) where the patient resides to facilitate specimen collection and appropriate follow-up. **To report a suspect measles case, contact the ECDOH Office of Epidemiology and Disease Surveillance at (716) 858-7697.**
 - Measles remains a common disease in many parts of the world and is introduced into the United States through international travel. Healthcare providers should ensure that all patients are up-to-date with age appropriate MMR (measles, mumps, rubella) vaccination or other evidence of measles immunity.

MEASLES EPIDEMIOLOGY

Measles can be severe and is highly infectious; following exposure, up to 90% of susceptible persons develop measles. It is spread by airborne contact with an infected person through coughing and sneezing. Measles virus can remain active and contagious for up to 2 hours in the air or on surfaces. The time from exposure to rash onset averages 14 days with a range of 7 to 21 days. Persons with measles are infectious from 4 days before to 4 days after rash onset.

CLINICAL FEATURES

Measles is characterized by a prodrome of fever (101–105 degrees F) followed by cough, coryza, and/or conjunctivitis. An erythematous, maculopapular rash presents 2-4 days later and lasts 5-6 days. It usually starts on the face and proceeds down the body to involve the extremities last and may include the palms and soles. The rash is usually discrete but may become confluent on the upper body; it resolves in the same order that it appeared. Koplik spots (punctate blue-white spots on the bright red background of the buccal mucosa) may be present, often before the rash develops, but are often not seen and are not required for the diagnosis of measles.

REPORTING DETAILS

Health care providers should increase their index of suspicion for measles in clinically compatible cases. Confirmed or suspected cases of measles are required to be reported immediately to the LHD. A report should be made at the time of initial clinical suspicion. The LHD should also be notified of discharge plans from the healthcare setting. This is especially important if the case lives in a multifamily dwelling, dormitory, group home or has young children at home.

INFECTION CONTROL

Measles is spread via airborne transmission and direct contact with infectious droplets. **Cases of fever and rash illness should immediately be placed in airborne isolation.** If an airborne infection isolation room is not available, then the exam room used to isolate a suspect measles case should not be used for 2 hours after the case leaves the room and the number of people entering and leaving should be minimized. When transporting a patient through the hospital, the patient should be masked. If possible, elevators and corridors should not be used for two hours after the patient has passed through them. If possible, any procedures required for the patient should be performed in the patient's room or delayed until the patient is no longer infectious. **If a suspect measles case, being evaluated as an outpatient, needs to be sent to a hospital emergency room, the emergency room should be notified ahead so that appropriate infection control precautions can be implemented upon arrival.**

LABORATORY TESTING

Viral specimens (throat or nasal-pharyngeal swab and urine) and serology (IgM and IgG) should be obtained for diagnostic testing and confirmation. Use of commercial laboratories for measles testing may take up to a week to obtain results. Reporting suspected cases of measles enables access to rapid testing through the NYS Wadsworth Center Laboratory. The LHD can assist in arranging testing at the Wadsworth Center Laboratory. Viral specimens that result in a positive PCR or culture will be forwarded to CDC for confirmation and genotyping.

MEASLES POST-EXPOSURE PROPHYLAXIS (PEP)

The successful initiation of measles PEP requires rapid intervention. The LHD can assist with the proper PEP recommendations and infection control measures. Measles vaccination should be administered to susceptible contacts of a measles patient within 72 hours of exposure and may offer protection. Immune globulin is indicated for susceptible household or other close contacts of patients with measles who are ineligible to receive MMR vaccination, particularly those contacts younger than 6 months of age, pregnant women and/or immunocompromised persons, for whom risk of complications is highest. Immune globulin should be given within 6 days of exposure to prevent or lessen the severity of measles.

MEASLES IMMUNITY

Acceptable presumptive evidence of immunity to measles includes:

- Born prior to 1957; or
- Written documentation of age-appropriate vaccination with 2 doses of measles-containing vaccine separated by at least 28 days for school-aged children (grades K-12) and adults at high risk for exposure and or transmission (i.e., healthcare personnel, students at post-high school educational institutions, and international travelers); or
- Written documentation of age-appropriate vaccination (i.e., aged ≥ 12 months) with at least 1 dose of measles-containing vaccine for preschool-aged children and adults who are not considered high risk; or
- Laboratory evidence of immunity; or
- Laboratory confirmation of disease.

MEASLES VACCINE RECOMMENDATIONS

Children ≥ 12 months, Adolescents, and Adults

- All children should receive an MMR vaccine at 12 – 15 months of age. The second dose of MMR is routinely administered at age 4 – 6 years typically before entering kindergarten but may be administered as soon as 28 days after the first dose.
Vaccination should be provided at the earliest opportunity based on the ACIP recommended schedule.
- Children over one year of age who have received one dose of MMR vaccine and who have been exposed in locations mentioned in this advisory should receive a second dose within 72 hours of exposure as long as 28 days have passed since the first dose.
- Children over one year of age and are planning travel outside the U.S. should receive a second dose as soon as possible as long as 28 days have passed since the first dose. Second doses of MMR are valid as long as they are administered after 12 months of age and at least 28 days after the first dose was administered.
- Anyone who lacks proof of measles immunity, as defined above, should receive at least one dose of MMR vaccine. Two appropriately spaced doses of MMR vaccine are recommended for health-care personnel, college students, and international travelers.

Children 6–11 months of age who are traveling outside the U.S.

- Should receive one dose of MMR vaccine prior to international travel.
- MMR vaccine given before 12 months of age should not be counted as part of the routine series. Children who receive MMR vaccine before age 12 months will need two more doses for a total of three doses, the first of which should be administered at 12 – 15 months of age and the second at least 28 days later (typically at age 4 – 6 years or before beginning kindergarten).

RESOURCES:

Contact Information:

- For questions or to report a suspect measles case, please contact the ECDOH Office of Epidemiology and Disease Surveillance at (716) 858-7697.

Additional Information:

- Frequently asked questions:
- www.erie.gov/measles
- Complete information on MMR vaccine recommendations:
www.cdc.gov/mmwr/pdf/rr/rr6204.pdf
- 2018 Immunization Schedules:
www.cdc.gov/vaccines/schedules/
- The NYSDOH Measles Fact Sheet is available at:
www.health.ny.gov/diseases/communicable/measles/fact_sheet.htm
- Destination specific travel immunization information is available on the CDC's Travelers' Health website at:

wwwnc.cdc.gov/travel/destinations/list

- For additional information on measles outbreak control measures, clinical presentation and diagnostic tests please refer to the CDC website at: www.cdc.gov/vaccines/pubs/surv-manual/chpt07-measles.html
- The NYSDOH Outbreak Control Manual is available at:
www.health.ny.gov/prevention/immunization/providers/outbreak_control_guidelines.htm
- CDC Measles Cases and Outbreaks:
www.cdc.gov/measles/cases-outbreaks.html
- CDC Measles Elimination:
www.cdc.gov/measles/about/faqs.html#measles-elimination
- Measles photos:
www.immunize.org/photos/measles-photos.asp

The ECDOH Health Alert & Advisory System is an e-mail notification system designed to alert community partners about important health related information. You can sign up to receive alerts & advisories at www.erie.gov/health/index.php?q=health-amp-human-service-professionals

Health Category Definitions:

Health Alert FLASH: conveys the highest level of importance due to a large-scale, catastrophic public health emergency; warrants immediate action or attention

Health Alert Priority: conveys the highest level of importance; warrants immediate action or attention to a health problem or situation

Health Advisory: provides important information for a specific incident or situation; may not require immediate action

Health Update: provides updated information regarding an incident or situation; no immediate action necessary

The Erie County Department of Health does not provide medical advice. The information provided on the Erie County Department of Health website is not an attempt to practice medicine and is not intended as a substitute for professional medical advice, diagnosis, or treatment. It is for informational purposes only. Always seek the advice of your personal physician or other qualified health provider with any questions you may have regarding a medical condition or issue. Never disregard professional medical advice or delay in seeking it because of the content found on the Erie County Department of Health website or this correspondence.

